

## Supplementary file 1

Table S1. Drug interactions of phenytoin

May increase concentration	May decrease concentration		
CYP450 enzyme inhibitors:	CYP450 enzyme inducers:	Protein binding:	Gastrointestinal absorption:
<ul style="list-style-type: none"> <li>• Acetazolamide</li> <li>• Allopurinol</li> <li>• Amiodarone</li> <li>• Amlodipine</li> <li>• Azapropazone</li> <li>• Capecitabine</li> <li>• Chloramphenicol</li> <li>• Chlorphenamine</li> <li>• Cimetidine</li> <li>• Clarithromycin</li> <li>• Clinafloxacin</li> <li>• Clobazam</li> <li>• Co-trimoxazole</li> <li>• Dicoumarol</li> <li>• Diltiazem</li> <li>• Disulfiram</li> <li>• Doxifluridine</li> <li>• Efavirenz</li> <li>• Eslicarbazepine acetate</li> <li>• Felbamate</li> <li>• Fenyramidol</li> <li>• Fluconazole</li> </ul>	<ul style="list-style-type: none"> <li>• Carbamazepine</li> <li>• Carboplatin</li> <li>• Ciprofloxacin</li> <li>• Cisplatin</li> <li>• Dexamethasone</li> <li>• Diazoxide</li> <li>• Folic acid</li> <li>• Hypericum perforatum</li> <li>• Loxapine</li> <li>• Nelfinavir</li> <li>• Nevirapine</li> <li>• Rifampicin</li> <li>• Theophylline</li> </ul>	<ul style="list-style-type: none"> <li>• Aspirin</li> <li>• Naproxen</li> <li>• Phenylbutazone</li> <li>• Salicylic acid</li> <li>• Some Sulphonamides</li> <li>• Tolbutamide</li> <li>• Valproate</li> </ul>	<ul style="list-style-type: none"> <li>• Some Nutritional Formulas</li> </ul>

<ul style="list-style-type: none"><li>• 5-Fluorouracil</li><li>• Fluoxetine</li><li>• Fluvoxamine</li><li>• Imipramine</li><li>• Isoniazid</li><li>• Itraconazole</li><li>• Methaqualone</li><li>• Methsuximide</li><li>• Miconazole</li><li>• Nortriptyline</li><li>• Omeprazole</li><li>• Oral Anticoagulants</li><li>• Oxcarbazepine</li><li>• Posaconazole</li><li>• Phenylbutazone</li><li>• Propoxyphene</li><li>• Risperidone</li><li>• Ritonavir</li><li>• Rufinamide</li><li>• Stiripentol</li><li>• Sulfadiazine</li><li>• Sulfamethizole</li><li>• Sulfamethoxazole</li><li>• Sulfaphenazole</li><li>• Sulfinpyrazone</li><li>• Sulthiame</li><li>• Tacrolimus</li><li>• Tamoxifen</li><li>• Ticlopidine</li><li>• Topiramate</li><li>• Trazodone</li><li>• Trimethoprim</li><li>• Valproic acid</li><li>• Verapamil</li></ul>			
---	--	--	--

<ul style="list-style-type: none"> <li>• Viloxazine</li> <li>• Voriconazole</li> <li>• Warfarin</li> </ul>			
(1, 3,4,5,6)	(1,3,4,5)	(1,2,3)	(1,4,7)
↖↑References↑↗			

## References

1. Nation RL, Evans AM, Milne RW. Pharmacokinetic drug interactions with phenytoin (Part I). *Clin Pharmacokin* 1990;18:37–60.
2. Neels HM, Sierens AC, Naelaerts K, et al. Therapeutic drug monitoring of old and newer anti-epileptic drugs. *Clin Chem Lab Med* 2004;42:1228–1255.
3. Patsalos PN, Spencer EP, Berry DJ. Therapeutic drug monitoring of antiepileptic drugs in epilepsy: A 2018 update. *Ther Drug Monit* 2018;40:526–548.
4. Lucas C, Donovan P. Medications:'Just a repeat': When drug monitoring is indicated. *Aust Family Phys* 2013;42:18–22.
5. Patsalos PN, Phenytoin. In: *Antiepileptic Drug Interactions: A Clinical Guide*, 3rd ed. London: Springer- Verlag, 2016:103-111 & 245-266.
6. Baxter K, Antiepileptics. In: *Stockley's Drug Interactions*. 9th ed. London: Pharmaceutical Press, 2010:592–662.
7. Williams NT. Medication administration through enteral feeding tubes. *Am J Health-Sys Pharm* 2008;65:2347–2357.